J. D. Baker & C. A. McGrath

AFCI/Gen IV Working Group Meeting Salt Lake City January, 2006



- INL is primarily involved with development and fabrication of actinide targets for neutron capture and neutron fission cross section measurements at LANSCE.
- In FY05 we developed electrodeposition techniques and fabricated targets of ^{239, 240, 242}Pu on 2 µm and 4 µm Ti foils for DANCE measurements and on large stainless foils for one-sided fission measurements.




- In FY 06 we are developing electrodeposition techniques to deposit actinide isotopes onto thin (6 µm), metallized, polymer films for improved DANCE and fission neutron cross section measurements.
- We fabricated and shipped to LANL two 252Cf calibration sources using the target holders designed for use in the PPAC detector.
- On a separately-funded project, we have received approval to begin work developing the reductive distillation or evaporations technique.




- The most recent target was ²⁴⁰Pu, DANCE target
 - 98.84% ²⁴⁰Pu, 0.735% ²³⁹Pu and 0.141% ²⁴¹Pu
 - Chemically purified to remove other actinide isotopes, ²⁴¹Am
 - Pu coated with ~100 μ g/cm² (~0.7 μ m) VYNS
 - 4 μm Ti, covered with 2 μm Ti
 - 0.887 mg ± 4.4 %















Measurement Capabilities

- Currently ≈ 5% accuracy
- Take targets back and dissolve to improve uncertainty.
- Working to develop NIST traceable calibrations for gamma-ray and alpha detectors on the order of 1-2%.
- With modeling and calibration we may be able to eliminate need to return targets.



- Future work -- FY06 and Beyond
 - Thin-film Pu-240 fission target
 - Pu-239, Pu-238 PPAC targets
 - Evaporated targets on thin films
- We acknowledge the Office of Basic Energy Sciences, U.S.
 Department of Energy, through the transplutonium element
 production facilities at the Oak Ridge National Laboratory for the
 use of (these isotopes)



